

Preliminary results from VLT (FORs1) photometry of NGC 6397

Andreuzzi G., Testa, V.

INAF-OAR, via Frascati 33, 00040 Monteporzio Catone (Italy)

Marconi G.

ESO-Chile, Casilla 19001, Santiago (Chile)

Alcaino G., Alvarado F.

Isaac Newton Institute, Casilla 8-9, Correo 9, Santiago (Chile)


Buonanno R.

University of Torvergata, 00100 Roma (Italy)

Abstract. We present VLT (FORs1) photometry of the lower main sequence (MS) of the Galactic Globular Cluster (GGC) NGC 6397, for stars located in 2 fields extending from a region near the cluster center out to $\simeq 10'$. The obtained CMD shows a narrow MS extending down to $V \simeq 27$ (figure c), much deeper than any previous ground based study and comparable with previous HST photometry (Cool et al. 1996). The comparison between observed MS Luminosity Functions (LFs) derived for 2 annuli at different radial distance from the center of the cluster shows a clear-cut correlation between their slope before reaching the turn-over, and the radial position of the observed fields inside the cluster area: the LFs become flatter with decreasing radius, a trend that is consistent with the interpretation of NGC 6397 as a dynamically relaxed system.

1. Introduction

GC stars fainter than the MSTO are substantially unevolved and are located along the MS according to their initial mass. Their luminosity distribution is usually described by the Initial Mass Function as modified by subsequent dynamical evolution to become the Present Day Mass Function. The mass segregation then should be seen as a radial dependence of the LFs. but to verify these theoretical predictions we need to obtain a statistically significative sample of data starting from the central regions of the cluster. Now this is possible also by ground observations thanks to the great power of the VLT and the FORs cameras.



2. Data analysis and discussion

Our observations consist of several 600 s exposures in the filter V and I Bessel (proposals 63.H-0721(A); 65.H-0531(B)) of two fields ($6.8' \times 6.8'$) partially overlapping and located at $\simeq 7.5'$ southeast of the center of the cluster. We have used HST observations (Cool et al., 1996 ApJ, 468, 655) in common with our fields (figure b) to make the calibration of the data. The LFs (in bin of 0.5 mag V) for the stars in the regions $R_1 < 5.5'$ and $5.5' \leq R_2 \leq 9.8'$ have been obtained by following the usual procedure and taking into account for the incompleteness of the images (figure a). It is evident from figure a) that both the observed LFs show a maximum at $V \simeq 22$, then decrease steadily to fainter luminosities maintaining a similar shape, but a tangible different slope. In particular the LFR2 ($19.5 \leq V \leq 22$), is steeper ($x = 0.16 \pm 0.01$) than the LFR1 ($x = 0.076 \pm 0.003$), suggesting that the relative ratio luminous stars/faint stars is greater within S1 than S2 and then confirming the mass-segregation effect in this cluster.¹

¹This work has been supported by the MURST/Cofin2000 under the project: Stellar observables of cosmological relevance.